

MOLECULAR
CELL

11
2003

QH
573

M654

V.11

2003

NO.5-6

LIFE

SCIENCE



THE LIBRARY
OF
THE UNIVERSITY
OF TEXAS
AT
AUSTIN

JOURNAL


APR 24 2004

Life Science Library MAI 220

Molecular Cell

Volume 11 Number 6

June 2003



**RNA
Recognition
by Smaug**

LIFE SCIENCE LIBRARY
MOLECULAR CELL
QH 573 N654v 11.6
7/3/03

Molecular Cell

Previews

- New Tricks for an Old Dogma:**
Riboswitches as *cis*-Only
Regulatory Systems G.D. Stormo 1419
- Enr1 Proteolysis:**
How SCF^{Enr1} Helps to Activate
the Anaphase-Promoting Complex J.-M. Peters 1420
- A New Link for a Linker Histone** A. Conconi and R.J. Wellinger 1421

Articles

- Substrate Specificity of Rhomboid Intramembrane
Proteases Is Governed by Helix-Breaking Residues
in the Substrate Transmembrane Domain** S. Urban and M. Freeman 1425
- Context of Multubiquitin Chain Attachment
Influences the Rate of Sic1 Degradation** M.D. Petruski and R.J. Deshaies 1435
- Structure of a β -TrCP1-Skp1- β -Catenin Complex:
Destruction Motif Binding and Lysine Specificity
of the SCF^{TrCP1} Ubiquitin Ligase** G. Wu, G. Xu, B.A. Schuman, P.D. Jeffrey,
J.W. Harper, and N.P. Pavletich 1445
- Insulin Activation of Rheb,
a Mediator of mTOR/S6K/4E-BP Signaling,
Is Inhibited by TSC1 and 2** A. Garami, F.J.T. Zwartkruis, T. Nobukuni,
M. Joaquin, M. Rocco, H. Stocker,
S.C. Kozma, E. Hafen, J.L. Bos, and G. Thomas 1457
- Multiple Roles of Tap42 in Mediating
Rapamycin-Induced Transcriptional Changes in Yeast** K. Düvel, A. Santhanam, S. Garrett,
L. Schreper, and J.R. Broach 1467
- JunD Mediates Survival Signaling
by the JNK Signal Transduction Pathway** J.A. Lamb, J.-J. Ventura, P. Hess, R.A. Flavell,
and R.J. Davis 1479
- Sustained Activation of the JNK Cascade
and Rapamycin-Induced Apoptosis
Are Suppressed by p53/p21^{ras}** S. Huang, L. Shu, M.B. Dilling, J. Easton,
P.C. Harwood, H. Ichijo, and P.J. Houghton 1491
- Identification of Transcription Factor KLF8 as a
Downstream Target of Focal Adhesion Kinase in Its
Regulation of Cyclin D1 and Cell Cycle Progression** J. Zhao, Z.C. Bian, K. Yee, B.P.C. Chen,
S. Chien, and J.-L. Guan 1503
- An Elongation Factor G-Induced Ribosome
Rearrangement Precedes tRNA-mRNA Translocation** A. Sevestarbergh, V.I. Katunin, D. Mohr, F. Peske,
M.V. Rodina, and W. Wintermeyer 1517
- TbMP57 Is a 3' Terminal Uridylate Transferase
(TUTase) of the *Trypanosoma brucei* Editosome** N.L. Ernst, B. Parisiucci, R.P. Igo, Jr.,
A.K. Panigrahi, R. Salavati, and K. Stuart 1525
- RNA Recognition via the SAM Domain of Smaug** J.B. Green, C.D. Gardner, R.P. Wharton,
and A.K. Aggarwal 1537
- Structure of an mRNA Capping Enzyme Bound
to the Phosphorylated Carboxy-Terminal
Domain of RNA Polymerase II** C. Fabrega, V. Shet, S. Shuman, and C.D. Lima 1549
- Modulation of NF- κ B Activity by Exchange of Dimers** S. Sacconi, S. Pantano, and G. Natoli 1563
- Structural Basis for Ligand-Independent Activation
of the Orphan Nuclear Receptor LHR-1** E.P. Sablin, I.N. Krylova, R.J. Fletterick,
and H.A. Ingraham 1575

(continued)

Modulation of NF- κ B Activity by Exchange of Dimers

Simona Saccani, Serafino Pantano,
and Giancchino Natoli*
Institute for Research in Biomedicine
Via Vela 6
CH6500, Bellinzona
Switzerland

Summary

Transcription factors within a family usually share the ability to recognize similar or identical consensus sites. For example, the five mammalian NF- κ B/Rel proteins generate more than 12 dimers recognizing 9–11 nucleotide κ B sites. Each dimer selectively regulates a few target promoters; however, several genes are redundantly induced by more than one dimer. Whether this property simply generates redundancy in target gene activation or underlies more complex regulatory mechanisms is an open issue. We show here that during dendritic cell maturation, rapidly activated dimers (e.g., p50/RelA) bound to a subset of target promoters are gradually replaced by slowly activated dimers (e.g., p52/RelB). Since the dimers have different transcriptional activity at each promoter, the dimer exchange allows fine tuning of the response over time. Further, due to the insensitivity of p52/RelB to the NF- κ B inhibitors, the I κ Bs, dimer exchange contributes to sustained activation of selected NF- κ B targets in spite of the resynthesis of I κ B α .

Introduction

The nuclear factor kappa B (NF- κ B) family of transcription factors (Sen and Baltimore, 1986) regulates numerous genes controlling immune response, cell growth, apoptosis, and tissue differentiation (Baldwin, 2001; Ghosh et al., 1996).

The five mammalian NF- κ B/Rel proteins contain an N-terminal segment of about 300 amino acids, the Rel-homology-domain (RHD), that is responsible for DNA binding, dimerization, nuclear translocation, interaction with the I κ Bs, and transcriptional regulation (Siebenlist et al., 1994; Verma et al., 1995). Three family members, p65 (RelA), cRel, and RelB, contain transcriptional activation domains (TAD) at the C terminus and therefore are able to directly activate transcription. The other two members, p50 and p52, are synthesized as large precursors (p105 and p100, respectively) with an N-terminal RHD and C-terminal ankyrin repeats; ubiquitin-dependent proteasomal processing removes the C-terminal domain and releases mature p50 and p52 (Botts and Nabel, 1996; Palombella et al., 1994; Xiao et al., 2001). p50 and p52 lack a TAD and therefore form homodimers with no intrinsic ability to activate transcription. However, they form transcriptionally active heterodimers in association with p65, cRel, and RelB. Moreover, p52

can activate transcription when complexed to Bcl-3, an I κ B-like molecule with coactivator functions (Siebenlist et al., 1994). Dimerization is required for NF- κ B binding to DNA, and more than 12 homo- and heterodimers have been described (Thanos and Maniatis, 1995; Verma et al., 1995). Different dimers are held in the cytoplasm by interaction with specific inhibitors: dimers containing p65 or cRel associate with the I κ Bs (inhibitors of κ B), which include I κ B α , I κ B β , and I κ B ϵ (Whiteside and Israel, 1997). I κ Bs contain an N-terminal regulatory region that is phosphorylated in response to stimulation (Brockman et al., 1995; Brown et al., 1995; DiDonato et al., 1996) and C-terminal ankyrin repeats, which mediate association with NF- κ B dimers (Hudford et al., 1998; Jacobs and Harrison, 1996). Conversely, RelB/p52 does not associate with the I κ Bs (Kistler et al., 1998; Lambeck et al., 1994; Wei et al., 1994) and is retained in the cytoplasm by p100 (Dobrzanski et al., 1999; Solan et al., 2002). I κ B-sequestered complexes are released by activation of the canonical NF- κ B activation pathway (Karin and Ben-Neriah, 2000), which depends on the IKK α /IKK β subunit of the I κ B-kinase (IKK) complex (DiDonato et al., 1997; Mercurio et al., 1997; Regnier et al., 1997; Woronicz et al., 1997; Zandi et al., 1997) and on its noncatalytic partner, IKK γ /NEMO (Mercurio et al., 1999; Rothwarf et al., 1998; Yamaoka et al., 1999). By phosphorylating two N-terminal serines in the I κ Bs, IKK β generates a docking site for the β TrCP proteins (Spencer et al., 1999; Winston et al., 1999; Yano et al., 1999), which polyubiquitinate the I κ Bs and target them for proteasomal degradation, thus liberating p65- and cRel-containing dimers. Release of p52/RelB (Siebenlist et al., 2001; Xiao et al., 2001) as well as p50/RelB dimers (Muller and Siebenlist, 2000) occurs through a “noncanonical” pathway requiring the NF- κ B-inducing kinase (NIK) (Xiao et al., 2001), which phosphorylates and activates IKK α (Regnier et al., 1997). In turn, IKK α phosphorylates p100, thus directing its polyubiquitination and processing (Siebenlist et al., 2001). This pathway is induced in response to a subset of stimuli such as BAFF, CD40 ligand, and LT β -R triggering (Claudio et al., 2002; Coope et al., 2002; Dejardin et al., 2002), and it has much slower activation kinetics than the canonical one. The independence of the two pathways is further indicated by the integrity of the noncanonical pathway in IKK α $-/-$ and IKK γ /NEMO $-/-$ cells (Dejardin et al., 2002), as well as by the observation that BAFF selectively activates the noncanonical pathway (Claudio et al., 2002). Whether the two pathways regulate completely distinct or partially overlapping sets of genes is still an open issue. Expression of several genes (e.g., *G-CSF*, *VCAM*, and *TNF α*) is elevated without stimulation in both IKK α $-/-$ mice (in which the major activated complex is p50/RelA) (Baj et al., 1995) and in mice lacking the C terminus of p100 but still expressing p52 (in which the major activated complex is p52/RelB) (Shikawa et al., 1997). On the other hand, analysis of gene expression induced by LT β -R triggering indicates that in fibroblasts, some genes are nonredundantly regulated by either of the two pathways (Dejardin et al., 2002).

An obvious requisite for redundancy is that more than

*Correspondence: giancchino.natoli@irb.unist.ch